

#### **DETOX & IMMUNE SUPPORT**

## **BIOTOXIN BINDER**

BioToxin Binder acts systemically, and contains a special form of carbon that has an enhanced binding ability. It also contains wild crafted **yucca root** and **extracts of fulvic acid**. It is specially formulated to support binding specific biotoxins like mycotoxins from mold and ammonia, as well as bacteria, parasites and other byproducts from yeast, fungi, and viruses.

Our BioToxin Binder is not formulated from activated charcoal. Activated charcoal is a long-chain carbon, which is classified as a "spent" carbon. What does this mean? Think of a spent carbon as a sponge that soaks up toxins only in the GI system. One of its weaknesses is that, just as a sponge, it can pick up and absorb, but it also can release (think "drip") things it has previously picked up along the way. This reality can lead to leaving things behind that more effective binders don't.


CellCore Biosciences products use long- and short-chain **BioActive Carbon** molecules, which have the ability to support life. The components that comprise BioToxin Binder include polyelectrolytes, polysaccharides, amino acids, and organic acids. This product works as a systemic binder as well as supporting cellular respiration (cellular energy production).

Our BioToxin Binder goes through a three-step proprietary process in which other molecules are attached to enhance its overall efficacy and ability to support the body as it removes biotoxins. Included in a comprehensive treatment protocol, BioToxin Binder is effective in supporting the body's efforts to remove the causes of toxicity and restore the body to overall systemic wellness.

Recommended dosage is two capsules twice daily, with or without food.



# **Supplement Facts**



Other Ingredients: HPMC (capsule)

**DAY 1 TO DAY 120** 

#### OPEN & DRAIN

### DAY 30 TO DAY 120

#### **BIND & REMOVE**

SUPPORT &





IN THE EVENT OF ADVERSE REACTIONS TO OUR PRODUCTS, REDUCE THE CURRENT DOSAGE. DO NOT COMBINE PARA 1 AND BINDERS. KEEP THEM AT LEAST AN HOUR APART FROM EACH OTHER. SEPARATE OUR PRODUCT FROM PRESCRIPTION DRUGS BY AT LEAST TWO HOURS. IT IS SAFE TO TAKE OTHER SUPPLEMENTS WITH OUR BINDERS, BUT AVOID TAKING THEM WITH PARA 1.

 $A void\ adding\ new\ protocols\ or\ supplement\ use\ if\ pregnant,\ breastfeeding\ or\ attempting\ to\ conceive.$ 

REV 10/2018